Zoologischer Anzeiger 255 (2015) 71…79

Contents lists available at ScienceDirect

Zoologischer Anzeiger

journal homepage: www.elsevier.com/locate/jcz

# Intercontinental island hopping: Colonization and speciation of the grasshopper genus Phaulacridium (Orthoptera: Acrididae) in Australasia

Julia Goldberg [a,], Mary Morgan-Richards [b], Steven A. Trewick [b]

[a] Department for Morphology, Systematics and Evolutionary Biology, J.F.Blumenbach Institute of Zoology & Anthropology, Georg-August-University Göttingen, Germany
[b] Ecology Group, Institute of Agriculture and Environment, Massey University, Private Bag 11-222, Palmerston North, New Zealand

article info

Article history:
Received 24 January 2014
Received in revised form 22 February 2015
Accepted 22 February 2015
Available

Table 1
DNA variation and haplotype diversity within and between regional samples of P. marginale in the New Zealand region and the two populations of P. otagoense, with the sample size for each region (n), number of observed haplotypes ($N_{haps}$), number of unique haplotypes ($N_{unihaps}$) per population, average number of nucleotide differences (k), nucleotide diversity ( × $10^{Š3}$), number of polymorphic sites (S), Tajima•s D (*P< 0.05) and haplotype diversity (h). Region abbreviations correspond to abbreviations in Supplementary Table 1 and Fig. 2B (NNZ = Northern North Island, CNZ = Central Mainland New Zealand, SNZ = Southern South Island, CHI = Chatham Islands; NZota Alex = P. otagoense population in Alexandra, Otago, NZota Mack = P. otagoense population in Mackenzie, Canterbury).

| Area | n | $N_{haps}$ | $N_{unihaps}$ | k | | S | Tajima•s D | h |
|---|---|---|---|---|---|---|---|---|
| NNZ | 16 | 7 | 4 | 0.358 | 0.81 | 2 | Š0.3817 | 0.342 |
| CNZ | 61 0 0 6.3761 499.0692 709.1268 Tm (Š)Tj /F1 1BT /F2 1 Tf .0002 0 0 -.0002 315.8074 753.9355 T 0 -.0002 339.97g4.3761 0 0 6.3761 169.8885 700.559 Tm (61 0 0 | | | | | | | |



Fig. 3. Mismatch distributions calculated for P. marginale (black) and P. otagoense (grey) from COI mtDNA illustrating the differences in genetic divergence within the two species. (A) Frequency distribution plot based on proportionated frequencies and pairwise genetic distances highlighting the difference in frequency distribution in the two species, with P. marginale mostly exhibiting low genetic distances (up to 1.5%) and P. otagoense showing the highest frequency at approximately 4% pairwise genetic distance. (B) Distribution plot based on genetic and geographic distances of individuals of the two species, depicting the high genetic divergence coupled with a narrow geographic range in P. otagoense (grey triangles) and the opposite …low genetic distance within a wider geographic distribution …in P. marginale (black dots).

networks, each representing grasshoppers from one of the two small areas in southern New Zealand where this species is found (Fig. 2A). This species displays higher genetic diversity than the widespread P. marginale with nine haplotypes found in a sample of 11 grasshoppers (Fig. 2A, Table 1). Of these nine haplotypes seven were unique within the respective population (Fig. 2B inset (i), Supplementary Table 1, Table 1). The calculated uncorrected pairwise genetic distances within this species reached 0.015 (1.5%) in the Alexandra population (NZotaAlex) and 0.018 (1.8%) in the more northern Mackenzie population (NZotaMack) (Fig. 2A). The maximal genetic distance between any two P. otagoense haplotypes was 0.037 (3.7%) across a spatial range of about 180 km. This is similar to the maximum genetic distance between P. otagoense and P. marginale (0.035/3.5%). Mismatch distributions calculated for P. marginale and P. otagoense highlights the difference in genetic divergence within these two species (Fig. 3A and B).

Populations of P. marginale displayed very low levels of DNA nucleotide diversity ( ) within mainland New Zealand and the Chatham Islands (Table 1). The sequences of the northern area had the highest level of haplotype and nucleotide diversity. These results of mtDNA differentiation refute the prediction of P. marginale partitioning into northern, southern and Chatham Island populations, but rather show panmixis in New Zealand. In the current data set no isolation by distance is apparent. Tajima D tests (Tajima, 1989) were only signi“cant for the SNZ dataset and the total population (Table 1). Negative Tajima•s D statistics for the different populations indicate an excess of sites with low frequency polymorphisms, i.e., the population has yet to reach equilibrium (Tajima, 1989). The two P. otagoense population samples on the other hand show higher estimates of than P. marginale populations in their small modern range (Table 1), implying that this species had a larger population size and range until relatively recently.

The Australian species P. vittatum cannot be


NZ
Aus
0.1
(B)
Chatham Is.
1
NZ, S.I.
1
P. marginale
NZ, N.I.
S.I., Otago, Mackenzie
1
P. otagoense
0.96
S.I., Otago, Alexandra
0.83
Tasmania
P. vittatum
1
Lord Howe Is.
P. howeanum
P. vittatum
1
Western Aus.
P. crassum
5 changes

the habitat/region occupied by P. otagoense today probably represents the most extensive natural lowland grass/scrubland available in New Zealand prior to human colonization. Other extensive

Vandergoes, M.J., Williams, P.W., NZ-INTIMATE members, 2007. Towards a climate event stratigraphy for New Zealand over

Sanmartin, I., Wanntorp, L., Winkworth, R.C.,2007. West wind drift revisited: testing for directional dispersal in the Southern Hemisphere using event-based tree “tting. J.Biogeogr. 34, 398…416.
Schmitt, T., 2007. Molecular