European Journal of Taxonomy 577: 1–46
https://doi.org/10.5852/ejt.2019.577
ISSN 2118-9773
www.europeanjournaloftaxonomy.eu
2019 · Hegg D.

This work is licensed under a Creative Commons Attribution Licence (CC BY 4.0).

Research article

urn:lsid:zoobank.org:pub:5ED633C5-4F9C-4F9D-9398-B936B9B3D951

Diversity and distribution of Pleioplectron + X W W R Q F D Y H Z W (Orthoptera: Rhaphidophoridae: Macropathinae), with the synonymy of Weta Chopard and the description of seven new species

Danilo HEGG[1], Mary MORGAN-RICHARDS[2] & Steven A. TREWICK[3,*]

[1] 135 Blacks Road, Opoho, Dunedin 9010, New Zealand.
[2,3] Wildlife & Ecology Group, School of Agriculture and Environment, Massey University, Private Bag 11-222, Palmerston North 4442, New Zealand.

[*] Corresponding author: s.trewick@massey.ac.nz
[1] Email: danilo_hegg@hotmail.com
[2] Email: M.Morgan-Richards@massey.ac.nz1 1

Abstract. The genus Thee I..aniaos15Tw Pleioplectronus T5.6D 5os15Tw<005A0044
EutrpepanJournial(ofT)92.12 axonomy https://doi.Mor/1

Introduction

7 K H V \ V W H P D W L F V R I 1 H Z = H D O D Q G F D Y H Z • W — D W W K H J H Q H the limited number of accessible and diagnostically informative morphological characters and often

Material and methods

Collection and morphological methods

&DYH Z•W— ZHUH FROOHFWHG RSSRUWXQLVWLFDOO\ DURXQG 1
and caves, and pitfall trapping. Two-letter codes in the Material examined sections below refer to the New Zealand entomological regions seen on Fig. 2C. More than 4300 specimens have been catalogued, and many have been examined in detail and sampled for DNA sequence comparison, including 378 specimens of Pleioplectron. Specimens are held in the Phoenix Lab collection at Massey University (MPN) with the exception of type material, which is lodged at Museum of New Zealand Te Papa 7RQJDUHZD 101= 5HOHYDQW VSHFLPHQV ZHUH LGHQWL¿HG E
Chopard (1923), Richards (1959) and Fitness et al. (2018).

Specimens were examined and photographed using a DSLR camera (Nikon D800, Nikon D850 or 6RQ\ . 5,, DWWDFKHG WR D 1LNRQ 3ODQ PLFURVFRSH WL
Cognisys Stackshot 3× automated rail. Focus stacks were generated using the software Helicon Focus 6.8.0 Pro (Helicon Soft Ltd, 2000). Adults were distinguished from immature individuals by darker, sclerotised bodies and fully formed external genital structures. In particular, the pigmentation, shape and sharpness of ovipositors, subgenital plates and cerci were informative about developmental stage. We looked for the presence/absence of each of 22 apical leg spines (Fitness et al. 2015) (Fig. 1), as well as the combinations and numbers of linear spines on the legs, the characteristics of the antennae and the shape of the subgenital and suranal plates. Measurements of key body parts were obtained using digital callipers (Table 1; also presented in Supplementary Material).

Collection acronyms

CMNZ = Canterbury Museum, Christchurch, New Zealand
iNaturalist = Available from iNaturalist.org [accessed 28 Apr. 2019]
LCR = Landcare Research, Wellington, New Zealand
01+1 0XVpXP QDWLRQDO GŒKLVWRLUH QDWXUHOOH 3DULV
MPN = Phoenix Lab, Massey University

Table 1. Dimensions and spine counts in species of *Plectroplectron* Hutton, 1896 (continued on next page).

		P. simplex	P. hudsoni	P. thomsoni comb. nov.	P. triquetrum sp. nov.	P. auratum sp. nov.
Sample size		, , ♂♂	, , ♂♂	, , ♂♂	, , ♂♂	, , ♂♂
Apical spines fore, mid and hind femora[1]		1 0 , 1 1 , 0 1	1 0 , 1 1 , 0 0	1 1 , 1 1 , 0 1	1 1 , 1 1 , 0 1	1 1 , 1 1 , 0 1
Apical spines fore, mid and hind tibiae		4-4-8	4-4-6	4-4-6	4-4-8	4-4-8
Body length (mm)[2]		14.6 (13.7–20.9)	9.4 (8.4–11.1)	14.7 (12.8–16.6)	11.2 (9.7–13.0)	11.55 (8.6–13.5)
Pronotum length (mm)		4.55 (3.8–5.4)	3.3 (2.8–3.6)	4.7 (4.2–5.2)	3.7 (3.2–4.2)	3.65 (3.1–4.9)
Ovipositor length (mm)		10.2 (9.9–11.7)	7.4 (6.8–8.1)	11.1 (10.2–12.4)	8.0 (7.6–9.0)	10.8 (9.9–12.3)
Ratio ovipositor to body length		0.71 (0.56–0.78)	0.78 (0.65–0.79)	0.78 (0.68–0.95)	0.73 (0.62–0.78)	0.96 (0.73–1.21)
Teeth: ventral valve of ovipositor		5 (4–6)	8 (7–9)	6 (5–6)	5 (4–6)	6 (5–7)
Teeth: dorsal valve of ovipositor		0	0	0	0	0
Length of hind tibia (mm)		15.0 (12.4–21.0)	11.2 (10.1–12.1)	20.0 (14.2–25.6)	14.0 (12.8–16.8)	14.3 (11.6–18.4)
Ratio hind tibia to body length		1.00 (0.84–1.33)	1.18 (1.05–1.35)	1.31 (0.97–1.74)	1.27 (1.09–1.49)	1.32 (0.99–1.57)
Superior spines hind tibia	prolateral retrolateral	26 (20–33) 28 (23–35)	20 (18–25) 21 (19–24)	40 (32–51) 41 (28–48)	30 (22–32) 27 (21–33)	33.5 (25–36) 32 (25–36)
Spine density on hind tibia (count/ mm)	prolateral retrolateral	1.61 (1.46–1.94) 1.87 (1.51–2.05)	1.79 (1.59–2.38) 1.96 (1.65–2.11)	2.12 (1.77–2.32) 1.98 (1.62–2.32)	2.05 (1.63–2.44) 1.89 (1.64–2.25)	2.21 (1.88–2.57) 2.24 (1.77–2.54)
Pairs of longer spines on hind tibia		0	0	0	0	1 (1–1)
Superior spines on 1st tarsus segment		8 (5–11)	8 (7–11)	12 (3–16)	9 (7–13)	9 (6–15)
Superior spines on 2nd tarsus segment		4 (0–5)	3 (2–4)	4 (1–7)	5 (3–9)	5 (2–7)
Fore tibia, inferior spines	prolateral retrolateral	3 (3–3) 3 (3–3)	1 (0–1) 2 (2–2)	2 (2–3) 2 (2–3)	2 (2–4) 3 (2–4)	3 (2–3) 3 (2–3)
Fore tibia, superior spines		0	0	0	0	0
Mid tibia, inferior spines	prolateral retrolateral	3 (3–3) 3 (2–3)	2 (1–3) 1 (0–2)	2 (1–2) 2 (2–3)	2 (1–3) 2 (1–3)	3 (2–3) 3 (1–3)
Mid tibia, superior spines	prolateral retrolateral	0	0	0	0	0
Fore femur, inferior spines	prolateral retrolateral	0	0	0	0	1 (0–2) 0
Mid femur, inferior spines	prolateral retrolateral	0	0	0	0	0
Hind femur, inferior spines	prolateral retrolateral	0 (0–1), v. small 3 (2–4), v. small	0 1 (0–1), v. small	0	0 5 (3–7), v. small	2 (0–2) 7 (4–8)

[1] The six numbers are, in order from left to right: fore femur prolateral and retrolateral, mid femur prolateral and retrolateral, hind femur prolateral and retrolateral. '1' means that an apical spine is present, '0' means that an apical spine is absent.

[2] Body length is measured from the apex of the fastigium to the posterior margin of the suranal plate.

Fig. 1.

(MPN CW1830), Miotopus diversus (Hutton, 1896) (GenBank accessions MK606169, MK606170, MK871353), M. richardsae (GenBank accessions MK532396, MK606171) and Pachyramma edwardsii (Scudder, 1869) (MPN CW746; GenBank accession MK871354) were used as an outgroup based on available phylogenetic information (Allegrucci et al. 2010; Fitness et al. 2018). A reduced taxon dataset with just 19 putative specimens of Pleioplectron allowed Bayesian inference to assess the monophyly of the genus. The Bayesian analysis used a chain length of 4 million generations sampling every 2000, and with a burn-in of 100000 generations. Maximum Likelihood was then used with mtDNA sequences from 88 putative specimens of



Fig. 2. Phylogenetic relationships of Pleioplectron + X W W R Q F D Y H Z • W — E D V H G R Q A. Bayesian analysis using a mtDNA sequence alignment of 1429 bp spanning the majority of COI, with a minimum individual sequence length of 1330. Posterior probabilities are shown by nodes. B. Maximum Likelihood tree using PhyML with a COI alignment that contained sequences ranging I U R P E S W R E S D O W K R X JE KS W DK WH W P D IM R • U LQ QaJ ZR IN WK H J H Q H Zealand entomological (Crosby) regions. These codes are indicated in the Material examined section for each species.

1RQH RI WKH FDYH Z•W— VSHFLPHQV H[DPLQHG FRXOG EH D

Hutton's original description of *P. hudsoni*, with three pairs of apical spines on the hind tibia, and so does the male subgenital plate, which is slightly longer than broad in a cuspidate shape and is slightly keeled. Here we consider that the widespread North Island morph represents *P. hudsoni*, but the brevity RI +XWWRQ¶V RULJLQDO GHVFULSWLRQ DQG WKH ODFN RI IHP the holotype specimen is missing (Richards 1959) and no other type material is available, we designate a male neotype for *P. hudsoni*.

Taxonomy

Order Orthoptera Latreille, 1793
Superfamily Rhaphidophoroidea Walker, 1869
Family Rhaphidophoridae Walker, 1869
Subfamily Macropathinae Karny, 1930
Tribe Macropathini Karny, 1930

Genus *Pleioplectron* Hutton, 1896

one anterior spine at the apex of the fore femur; four out of ten species additionally have a posterior spine at the apex of the fore femur (Table 1). All species have both an anterior and a posterior spine at the DSH[RI WKH PLG IHPXU 7KH KLQG IHPXU KDV D UHWURODWHUD All species have four apical spines at the apex of both fore and mid tibia, and at the apex of the hind tibia a pair of dorsal apical spines, a pair of dorsal sub-apical spines and a pair of ventral apical spines. Four out of ten species additionally have a pair of ventral sub-apical spines at the apex of the hind tibia (Table 1). The dorsal apical spines on the hind tibia are very long, generally at least twice or three times the length of ventral apical spines. Linear spines: fore and mid femur generally unarmed, except in Pleiplectron auratum sp. nov. and P. rodmorrisi sp. nov. Hind femur generally armed below; the number of spines varies both within and between species. Fore tibia always unarmed above; mid tibia generally unarmed above, except in Pleioplectron rodmorrisi sp. nov. Fore and mid tibia armed below, with one to three pairs of linear spines; the number varies both within and between species. The hind tibia is always armed above with two parallel rows of 20 or more (up to 50) linear spines; P. crystallae sp. nov. has fewer than 20 dorsal spines in each row (Table 1). Linear spines on the hind tibiae are socketed but not articulated, very variable in size on the same animal, randomly smaller and larger without obvious

Fig. 4. +HDG RI FDYH Z • WPleioplectron Hutton, 1896, showing sexual dimorphism. A B. P. simplex Hutton, 1896. A $GXOW ƒ +LQHZDL 5HVHUYH %DQNV 3H B $GXOW , +HOLFRSWHU +LOO 7UDFN P. crystallae sp. nov. Adult 031 &: ƒ %UDQFK &UHHN +XW 0WD2 ZSHQX OW31, &6DOLVEXU\ /RGJH 0W $U (MPN CW3941). Scale bar = 5 mm.

pattern (Fig. 7). Additionally, two species (*B. auratum* sp. nov. and *B. crystallae* sp. nov.) have one or more (up to four) pairs of much larger, possibly articulated dorsal spines on the hind tibiae. Fore and mid tarsi: all segments unarmed. Hind tarsi: always armed above with a variable number of small spines

2012; B. Taylor-Smith and M. Morgan-Richards leg.; leaf litter, forest fragment; night search; MPN CW2137, CW2186 to CW2188, CW2201, CW2403, CW2418, CW2463 ƒ , , Waimakariri 5 L Y H U & D V V ¿ H O G V W D W L R Q ƒ 6 ƒ (P D V O overhang near stream; MPN CW327A, CW327B, CW328, CW329, & U D L J L H E X U Q + H O L F R Track; 43.14628° S, 171.72807° E; 800 m a.s.l.; 19 Mar. 2006; P.M. Johns and R. Pratt leg.; in native forest; pitfall trap; MPN CW359 ‡ , , same locality as preceding; 10 Feb. 2017; D. Hegg leg.; clay bank on side of walking track; night search; MPN CW3459, CW3914, 3 R U W H U V 3 D V V) F Peak; 43.29361° S, 171.74232° E; 1000 m a.s.l.; 19 Mar. 2006; P.M. Johns and R. Pratt leg.; near stream; pitfall trap; MPN CW361 ‡ , , ƒ Lake Coleridge, H.E. Hart Arboretum; 43.36282° S, 171.53006° E; 400 m a.s.l.; 18 Nov. 2016; D. Hegg leg.; exotic forest; night search + insect net; MPN CW3178 to CW3181 • , Q \ P S K 0 W 6 R P H U V 6 K D U S O L Q) D O O V a.s.l.; 13 Oct. 2017; D. Hegg leg.; under boulder in native forest; night search + insect net; MPN CW3786, CW4001. Dunedin (DN) ‡ ƒ ƒ , Long Beach; 45.7444° S, 170.6406° E; 0 m a.s.l.; 10 May 2016; D. Hegg leg.; in sea cave; night search + insect net; MPN CW3009 to CW3013; same collection data as for preceding; 8 Nov. 2016; MPN CW4013 ƒ Opoho; 45.85399° S, 170.5339° E; 120 m a.s.l.; 14 Feb. 2016; D. Hegg leg.; in wooden retaining wall on roadside; night search + insect net; MPN CW2821 ‡ ƒ Q \ P S K Otago Boys' High School; 45.87083° S, 170.49588° E; 100 m a.s.

Originally described by Hutton (1896), the species was redescribed adequately by Richards (1959). A notable detail omitted by both Hutton (1896) and Richards (1959) is the presence of sensory hair tuffs on the antennal segments of some, but not all, sexually mature males (Fig. 5).

Male *Pleioplectron simplex* on Banks Peninsula tend to reach larger dimensions than elsewhere, but our JHQHWLF GDWD SURYLGH QR HYLGHQFH WKDW WKH %DQNV 3HQ measured from Banks Peninsula, the smallest (MPN CW4112) measured 16.3 mm in body length, 12% longer than the largest adult male we collected elsewhere (MPN CW4013 from Long Beach, Dunedin, 14.5 mm long). To avoid introducing geographical bias in our measurements, we only included one adult male *P. simplex* from Banks Peninsula in Table 1, which gives us an estimate for the maximum size of the species.

Pleioplectron hudsoni Hutton, 1896
Figs 2A–B, 6G, 7B, 8D–F, 10D–F, 14E, 18A, 19

Pleioplectron hudsoni + X W W R Q S O ¿ J V D

Diagnosis

$ VPDOO EURZQ FDYH Z•W— ZLWK D \HOORZ PHGLDQ GRUVDO V North Island, New Zealand. It could be most easily confused with *Miotopus diversus* (Hutton, 1896) or *Neonetus variegatus* Brunner von Wattenwyl, 1888, two species with which it is sympatric and shares the forest habitat. From both of these species it is distinguished by the lack of a posterior apical spine on the fore femur. Additionally, it is smaller than *M. diversus* and lacks dorsal linear spines on the mid tibia.

Etymology

Named after the New Zealand entomologist George V. Hudson (1867–1946), who collected the holotype specimen (a male) in Wellington (Richards 1959). According to Richards (1959), the holotype is now missing.

Material examined (see also Table 3 in Supplementary Material)

Neotype

1 (: = ($ / $ 1 ' ‡ ƒ D G X O W : H O O L Q J W R Q : / = H D O D Q G L D (F R V D Q F W 250 m a.s.l.; 25 Feb. 2017; Edward (Ted) Trewick, Mary Morgan-Richards and Steve Trewick leg.; in native forest; NMNZ AI.037500 (previously MPN CW3561).

Representative female

174.92389° E; 50 m a.s.l.; 21 Oct. 1999; S. Trewick leg.; under rocks at base of beech tree; MPN CW171A, CW171B ‡ ƒ , Kapiti Island; 40.85336° S, 174.92955° E; 100 m a.s.l.; 14 Nov. 2011; B. Taylor-Smith and S. Trewick leg.; MPN CW1993, CW1995 ‡ ƒ ƒ Tararua Forest, Otaki Forks; 40.88085° S, 175.21887° E; 150 m a.s.l.; 4 Nov. 2017; D. Hegg leg.; in roots of uprooted tree; night search + insect net; MPN CW3889, CW3890 , Levin, Lake Papaitonga; 40.6444° S, 175.23394° E; 10 m a.s.l.; 20 Sep. 2004; S. Trewick leg.; MPN CW161 • 1 nymph; Tararua Forest, Makahika Stream; 40.63057° S, 175.41321° E; 200 m a.s.l.; 20 Mar. 2004; M. Morgan-Richards leg.; MPN CW171 , 3 D O P H U V W R Q 1 R U W K 7 X U L W H D 5 R D G ƒ 6 ƒ (leg.; on ground under gorse bush; MPN CW186 ƒ same locality as preceding; 18 Nov. 2004; M. Morgan-Richards and S. Trewick leg.; under porch of house; MPN CW187 ƒ ƒ Palmerston North, Turitea Reserve; 40.43039° S, 175.67276° E; 150 m a.s.l.; 2008; M. Morgan-Richards and S. Trewick leg.; MPN TD17(1)A, TD19(1)A ‡ ƒ ƒ , same locality as preceding; 15 Aug. 2010; S. Trewick leg.; near boat shed; MPN CW1440 to CW1442 ƒ ƒ , , same locality as preceding; 19 Nov. 2010; S. Trewick leg.; night search; MPN CW1454, CW1457, CW1464, CW1465 , Palmerston North, Esplanade; 40.37222° S, 175.61740° E; 25 m a.s.l.; Apr. 2006; J. Goldberg leg.; MPN CW434 , 0 D Q D Z D W X * R U J H ƒ 6 ƒ (P D V O $ S U CW441. – Rangitikei (RI) • 1 ƒ Pohangina River, Totara Reserve; 40.12239° S, 175.85601° E; 150 m a.s.l.; 22 Aug. 2004; E. Trewick and S. Trewick leg.; MPN CW131 , Q \ P S K 5 X D K L Q H Forest, Alice Nash Memorial Heritage Lodge; 39.95591° S, 176.01522° E; 700 m a.s.l.; 28 Oct. 2018; D. Hegg leg.; on clay bank on side of track; night search + insect net; MPN CW4134, CW4140 Ruahine Forest, Gold Creek Track; 39.76367° S, 176.19614° E; 700 m a.s.l.; 25 Mar. 2014; M. Lusk leg.; in standing dead wood; MPN CW2678 ƒ , Ruahine Forest, Parks Peak Track; 39.68° S, 176.25° E; 9 Mar. 2012; M. Lusk leg.; MPN CW1916, CW1917 ƒ Rangitikei River, Mangaweka campsite; 39.81084° S, 175.80666° E; 300 m a.s.l.; 11 Sep. 2004; E. Trewick and S. Trewick leg.; in decaying log; MPN CW143. – Wanganui (WI) ‡ , , Bushy Park; 39.79690° S, 174.93084° E; 250 m a.s.l.; 29 Dec. 2013; M. Morgan-Richards and S. Trewick leg.; night search; MPN CW2496, CW2499. – Taranaki (TK) ‡ ƒ , , Lake Rotokare; 39.45409° S, 174.40923° E; 200 m a.s.l.; 2008; J. Fitness leg.; pitfall trap; LCR LR-L5-b, LR17-live-a, LR19(6)-live. – Wairarapa (WA) • 1 ƒ Bideford; 40.86653° S, 175.86795° E; 150 m a.s.l.; 18 Sep. 2004; M. Morgan-Richards and S. Trewick leg.; under rotting logs; MPN CW157 ‡ ƒ Puketoi Range, Towai Road; 40.41614° S, 176.14002° E; 600 m a.s.l.; 22 Oct. 1999; R. Brown leg.; MPN CW85. – Hawkes Bay (HB) • 1 ƒ Elsthorpe Reserve; 39.91961° S, 176.81603° E; 150 m a.s.l.; 29 Dec. 2005; S. Trewick leg.; MPN CW213 ƒ Mohi Bush; 39.85725° S, 176.90068° E; 450 m a.s.l.; 27 Nov. 1995; M. Morgan-Richards and S. Trewick leg.; inside log; MPN CW601 ‡ ƒ , , same locality as preceding; 1 Jan. 2012; M. Morgan-Richards and S. Trewick leg.; MPN CW1755, CW1759, CW1762 , same locality as preceding; 8 Oct. 2012; M. Morgan-Richards and S. Trewick leg.; MPN CW1861 , Q \ P S K Waipatiki Reserve; 39.28256° S, 176.96400° E; 100 m a.s.l.; 27 Jun. 2011; M. Lusk leg.; MPN CW1972, CW1973 , Kaweka Range, Smith-Russell Track; 39.37815° S, 176.33550° E; 600 m a.s.l.; 4 Feb. 2014; M. Lusk leg.; under log; MPN CW2680 ‡ , Kaweka Range, Kuripapango Lakes; 39.36148° S, 176.36078° E; 700 m a.s.l.; 16 Feb. 2014; M. Lusk leg.; under log; MPN CW2682 ƒ Kaweka Range, Makahu; 39.2184° S, 176.48407° E; 460 m a.s.l.; 6 Jan. 2014; M. Lusk leg.; under log; MPN CW2692 ƒ Maungahuru Range, Maungahuru Track; 39.20474° S, 176.68408° E; 750 m a.s.l.; 10 May 2011; M. Lusk leg.; MPN CW1609. – Bay of Plenty (BP) ‡ , Manganuku Campsite, Waioeka River; 38.29018° S, 177.38637° E; 200 m a.s.l.; 12 Apr. 1998; S. Trewick leg.; MPN CW561A ƒ same locality as preceding; 2 Jan. 2011; M. Morgan-Richards leg.; MPN CW1984 ‡ ƒ



Fig. 6. ' R U V D O Y L H Z V R I F D Y H O H U L D Q V S H F L P H Q V Z L W K H D G X O W , + H O L F R S W H U + L O O 7 U D F N . D L N R Q Y D G X O W ƒ & O D U H P R Q W % X V K G R U V D O Y L H Z V R I F D Y H O H U L D Q V S H F L P H Q V Z L W K H D G X O W ƒ : K L W H V % H D F K r o d r i g o , 5 D L O Z D \ 7 U D F N . D L N R Q Y D G X O W ƒ + R G G H U 5 L Y H U , Q O D Q G G D P L Q À X H Q F H W U D F N 0 3 1 & D G X O W Karori, Wellington (MPN CW3564). H. caudatum V S Q R Y D G X O W ƒ % U H Z V W H U + X (MPN CW4049). I. 3 À D Y L V R U Q R Y D G X O W ƒ % U H Z V W H U + X W 7 U D F N J. P. crystallae V S Q R Y D G X O W , & D Y H % U R R N * R X O D Q G ' R Z Q V 0 3 1

HEAD. Frons dark brown, covered with very sparse setae; vertex glabrous and of variegated colour. Eyes dark, bulging and large compared to size of head. Fastigium dark, with a white spot on either side.

Diagnosis

$ PHGLXP VL]HG FDYH Z•W— ZLWK ORQJ OHJV SDOH FRORXU is common and widespread in caves and forests in South Canterbury and Otago, New Zealand. It has prolateral and retrolateral apical spines on both fore and mid femur.

Etymology

Named in honour of Dr. J. Allan Thomson (1881–1928), director of the Dominion Museum, Wellington, who sent the type specimens to Chopard. It is not known whether Thomson collected the specimens

CW3928, CW3929, CW3970. North Canterbury (NC) ‡ *f f* , , Q \ P S K; same collection

point; relatively short (approximately three quarters of body length). Upper valve smooth above; lower valve with 4 to 6 strong teeth at apex on ventral edge (Fig. 10K–L). Dorsal view of terminalia as in P. simplex (Fig. 12A).



Fig. 7. / H I W K L Q G W L E L D R I P l e i o p l e c t r o n w u t t o n i , 1 9 9 6 , d o r s a l v i e w o f t e r m i n a l i a (Q R Y
+ X W W R Q D G X O W , + L Q H Z D L 5 H V H U Y H + X W W R Q 0 3 1 & G X O W
Otaki Forks, Tararua Forest (MPN CW3890). P. thomsoni & K R S D U G F R P E Q R Y
Claremont Bush Reserve, South Canterbury (MPN CW4115). P. triquetrum V S Q R Y D G X O W ƒ +
7 U D F N . D L N À X U D H 3 1 a & a t u m V S Q R Y D G X O W ƒ : K L W H V % H D F
CW4295). F. P. gubernator V S Q R Y D G X O W ƒ / H Z L V G . 3 . D u t a t u m s 1 n o & ;
D G X O W ƒ 0 R H U D N L 5 L Y H U À 3 1 L V R U Q R Y D G X O W , % U H Z V W H U +
Pass (MPN CW3896). P. crystallae V S Q R Y D G X O W ƒ & D Y H % U R R N * R X O D Q G
J. P. rodmorrisi V S Q R Y D G X O W ƒ + R G G H U 5 L Y H U , Q O D Q G . D L N À X U H
5 mm.

Pleioplectron auratum sp. nov.
urn:lsid:zoobank.org:act:97429B5F-54CF-4891-8174-64F038866150
Figs 2A–B, 6E, 7E, 8M–O, 10M–O, 15A, 18E

Diagnosis

$ VPDOO WR PHGLXP VL]HG FDYH Z•W— IRXQG LQ IRUHVWV DQG corner of South Island. The habitus is typical of Pleioplectron, chequered grey-brown in colour, with visibly banded legs and with a dorsal median line (Fig. 15A). A prominent pair of longer spines stands out on the hind tibia, about one third of the way up from the apex (Fig. 7E). Relative to body size, the ovipositor is longer than in any other species of this genus. Pleioplectron auratum sp. nov. is most similar to P. triquetrum sp. nov., although it i EMC /S [(P)129.0590041999 >>BDCf /Fm1 DhC /S [(P(MC /SpEBD



Fig. 8. $ G X O W P D O H W H U P L Q D O L P D J H R I Pleioplectron Hutton, 1896. Left column: ventral view (subgenital plate); central column: dorsal view; right column: lateral view. P. simplex Hutton, 1896, Hinewai Reserve, Banks Peninsula. A B. MPN CW4109. C. MPN CW4118. P. hudsoni Hutton, 1896, Otaki Forks, Tararua Forest Park (MPN CW3890). P. thomsoni (Chopard, 1923) comb. nov., Trotters Gorge, Otago (MPN CW3780). J-L. triquetrum sp. nov. J, L. MPN CW3922. K. MPN CW3928. M O. P. auratum sp. nov. M, O. Whites Beach, Rarangi (MPN CW4295). N. Waihopai River, Blenheim (MPN CW4146). Scale bars = 1 mm.

Legs. Moderately long; hind femora up to 20% longer than body; hind tibiae up to 50% longer than body in both males and females. Coxae and trochanters mottled light and dark brown. Fore and mid femora and tibiae with conspicuous alternating light and dark bands. Hind femora variegated; hind

Etymology

‹ * Î E H U Q — W Â U Œ L V / D W L Q I R U µ K H O P V P D Q ¶ E H F D X V H R I W K H Y I (Fig. 9A, C), which resembles the rudder under a boat.

Material examined (see also Table 7 in Supplementary Material)

Holotype

1 (: = ($ / $ 1 ' ‡ ƒ D G X O W % X O O H U % 5 / H Z L V 3 D V V & D Q Q L E D O * 840 m a.s.l.; 3 Feb. 2017; Danilo Hegg leg.; in rotting tree stump in native forest; night search + insect net; NMNZ AI.037484 (prev. MPN CW4063).

Paratype

Female terminalia. Subgenital plate tri-lobed, with middle lobe pointed, two outer lobes broad and rounded (Fig. 11A). Ovipositor reddish brown, recurved above near tip, terminating in a very sharp point; approximately 80% of body length. Upper valve smooth above; lower valve with 8 to 12 small but strong teeth at apex on ventral edge (Fig. 11B–C). Dorsal view of terminalia with a sharp point on last tergite, as in *P. crystallae* sp. nov. (Fig. 12B).

Pleioplectron caudatum sp. nov.
urn:lsid:zoobank.org:act:7ADDB012-690F-4A3F-83F9-FDC54F8F9EED
Figs 2A–B, 6H, 7G, 9D–F, 11D–F, 12C, 15C, 18G

Diagnosis

Head. Frons dark brown, with very sparse setae; vertex glabrous and of variegated colour. Eyes dark and bulging. Fastigium dark, with a white spot on either side. Scapes and peduncles pale and pilose; DQWHQQDH GDUN EURZQ FRYHUHG LQ ¿QH VHWDH 0D[LOODUL



Fig. 10. $ G X O W I H P D O H W H U P L Q D O L Pleioplectron Hutton, 1896. Left column: subgenital plate; central and right columns: oviposA C. P. simplex Hutton, 1896, Hinewai Reserve, Banks Peninsula (MPN CW4114). D F. P. hudsoni Hutton, 1896, Karori Ecosanctuary, Wellington. D. MPN CW3554.E F. MPN CW3552. G I. G. Pioneer ark, Opihi River, South Canterbury (MAPN CW3913). L. Trotters Gorge, Otago (MPN CW4007). L. M Q uratum N CW4120. N O. MPN CW4119. Scale bars = 1 mm.

Thorax. Pronotum variegated brown, mainly dark, with a few pale patches; dark with pale bars next WR SRVWHULRU PDUJLQ FRYHUHG LQ VSDUVH ¿QH WRPHQWX posterior margin nearly straight.

Legs. Moderately long; hind femora of approximately equal length as body; hind tibiae on average 20% longer than body in both males and females. Coxae and trochanters mottled light and dark brown. Fore and mid femora and tibiae with alternating conspicuous light and dark bands; hind legs variegated. Fore coxae with a pronounced lateral spine. Fore femora without linear spines above or below; armed with one prolateral spine at apex. Fore tibiae armed below, generally with one (at most two) linear spines on anterior edge, two linear spines on posterior edge. Fore tibiae armed with two long spines below (one prolateral and one retrolateral) and two short spines above (one prolateral and one retrolateral) at apex. Mid femora without linear spines above or below, but armed with one prolateral and one retrolateral spine at apex. Mid tibiae with two linear spines below on both anterior and posterior edge. Mid tibiae

Etymology

‹) O — Y LQ ERLVQ LQ W L Q IRU µ\HOORZ KRUQHG¶ LQ UHIHUHQFH WR (Figs 6I, 15D).

Fig. 11. $GXOW IHPDOH WHUPLQDOLD Pleioplectron Hutton, 1896. Left column: subgenital plate; central and right columns: ovipositor. A–C P. gubernator sp. nov. A. Awapoto Hut, Abel Tasman NP (MPN CW4057). B. C. Lewis Pass (MPN CW4071). D–F. P. caudatum sp. nov., Brewster Hut Track, Haast Pass (MPN CW4040). G–I. P. crystallae sp. nov., Brewster Hut Track, Haast Pass (MPN CW4042). J–L. P. crystallae

Material examined (see also Table 9 in Supplementary Material)

Holotype
1 (: = ($ / $ 1 ' ‡ ƒ D G X O W : H V W O D Q G : ' + D D V W 3 D V V % U H Z V
ƒ (P D V O 1 R Y ' D Q L O R + H J J O H J R Q I R U H
NMNZ AI.037488 (prev. MPN CW4055).

Paratype
1 (: = ($ / $ 1 ' ‡ , D G X O W V D P H F R O O H F W L R Q G D W D D V I R U K R O
CW4042).

Other material
NEW ZEALAND – Westland (WD) ‡ ; Haast Pass, Brewster Hut Track; 44.08062° S, 169.39055° E;
P D V O) H E ' + H J J O H J R Q I R U H V W À R R U ƒ Q L J K W V
, , Q \ P S K; same collection data as for holotype; MPN CW4041, CW4043 to CW4048, CW4050,
CW4051, CW4054 ‡ , , V X E D G X O W V 0 R H U D N L 5 L Y H U % O R Z À \ + X W
D V O ' H F ' + H J J O H J R Q I R U H V W À R R U Q L J K W V H D U
Otago Lakes (OL) • 1 nymph; Haast Pass, Bridle Track; 44.12142° S, 169.34396° E; 460 m a.s.l.;
13 Oct. 2018; D. Hegg leg.; on bank on side of track; night search + insect net; MPN CW4122.

Description

Measurements. See Table 1.

Head. Frons dark brown except for two white lateral patches; with very sparse setae; vertex glabrous and of variegated colour. Eyes dark and bulging. Fastigium dark, with a white spot on either side. Scapes and

Legs. Moderately long; hind femora of approximately equal length as body; hind tibiae on average 15% longer than body in both males and females. Coxae and trochanters pale with dark patches. Fore and mid femora and tibiae with conspicuous alternating light and dark bands, and hind legs variegated. Fore coxae with a pronounced lateral spine. Fore femora without linear spines above or below, but armed with one prolateral spine at apex. Fore tibiae armed below, generally with two linear spines on both anterior and posterior edge. Fore tibiae armed with two long spines below (one prolateral and one retrolateral) and two short superior spines (one prolateral and one retrolateral) at apex. Mid femora without linear spines above or below, but armed with one prolateral and one retrolateral spine at apex. Mid tibiae with up to two linear spines below on both anterior and posterior edge. Mid tibiae armed with two long spines below (one prolateral and one retrolateral) and two short spines above (one prolateral and one retrolateral) at apex. Hind femora unarmed below and at apex. Hind tibiae armed with 23 to 30 linear spines above, of varying length, on both anterior and posterior edges. Hind tibiae with two superior subapical spines (one prolateral and one retrolateral), two superior apical spines (one prolateral and one retrolateral) and two inferior apical spines (one prolateral and one retrolateral). Superior subapical spines and inferior apical spines of approximately equal length, superior apical spines approximately twice as ORQJ +LQG WDUVL ZLWK IRXU VHJPHQWV ¿UVW DQG VHFRQG VHJPHQW ZLWK ± VPDOO GRUVDO DOWHUQDWH VSLQHV 6HF VSLQHV 6HFRQG DQG IRXUWK VHJPHQWV RQH WKLUG WR RQH third of length of second and fourth segments.

Abdomen. Colour variegated brown, mostly reddish brown with black patches and covered in sparse, ¿QH WRPHQWXP 'RUVDO \HOORZ PHGLDQ OLQH WKLQ EXW JHQ

Male terminalia. Cerci long, pointed at apex, brown in colour and clothed in setae. Styli short and not visible from above. Subgenital plate broad, oval (Fig. 9G). Suranal plate a tall truncated cone, almost half length of cerci, and covering styli and all genital organs entirely (Fig. 9H).

Female terminalia. Subgenital plate tri-lobed, all lobes shallow and rounded (Fig. 11G). Ovipositor reddish brown, weakly recurved above at tip and relatively short (approximately three quarters of body length). Upper valve smooth above, but lower valve with 9 to 11 shallow teeth on ventral edge at apex (Fig. 11H–I). Terminalia similar to those of *P. simplex* when viewed from above (Fig. 12A).



Fig. 13. *Pleioplectron thomsoni* & K R S D U G F R PA E Original drawing by Luke Wen*f* & K R S D U G V \ Q W \ S H *f* 5 D L Q F O L I I 5 H V H U Y H B. Pixner & D Q W H Park, South Canterbury (MPN CW3912). Scale bar = 2 mm.

Legs. Relatively short; hind femora generally shorter than body length. Hind tibiae up to 15% longer

THORAX. Pronotum pink orange, with a prominent yellow median line, and black regions next to all PDUJLQV FRYHUHG LQ VSDUVH ¿QH WRPHQWXP $OO PDUJLQV rounded, posterior corners more angular.

LEGS. Very long, sexually dimorphic (see Fig. 17), longer in males than in females. Coxae and trochanters pale pink/orange. Fore and mid femora orange. Hind femora variegated, purple in proximal portion and



Fig. 17. Sexual dimorphism in the legs of Pleioplectron rodmorrisi sp. nov., with boxplot for a sample of 8 adult males and 7 adult females. 'Leg length' was calculated as the sum of the lengths of femur and tibia. A B. Photographs of live specimens of P. rodmorrisi sp. nov. in their natural environment. Kahutara Saddle, Seaward Kaikoura Range. $ G X OW $ GX OW. Box plot for the ratio of leg length/body OHQJWK E\ VH[$ RQH ZD\ 129 WHVW RQ OHJ OHQJWK E\ VH[in males than they are in females, with the following results:
Fore leg: F = 198.5 on 1 and 13 degrees of freedom; P = 10-9; R^2 = 0.9385.
Mid leg: F = 130.9 on 1 and 13 degrees of freedom; P = 10-8; R^2 = 0.9097.
Hind leg: F = 16.18 on 1 and 13 degrees of freedom; P = 0.0010; R^2 = 0.5545.

Pleioplectron simplex

Differentiation of Pleioplectron from other genera of Rhaphidophoridae in New Zealand

In the generic description we listed traits that are characteristic of Pleioplectron. Many of them, however, are not unique to Pleioplectron, and are not necessarily useful to differentiate it from other genera of New Zealand Rhaphidophoridae.

:DUG PDGH D ¿UVW DWWHPSW DW FRPSLOLQJ D NH\ WR 1H included the following features: apical spine count on fore, mid and hind femora; apical spine count on fore, mid and hind tibiae; shape of male subgenital plate; shape of female subgenital plate. Johns & Cook (2013) included a table as an appendix to their generic description of Wanoweta, in which they compared six genera of New Zealand Rhaphidophoridae based on the following traits: body length; apical spine count on fore, mid and hind femora; ventral linear spines on hind femora; dorsal linear spines on hind tibiae; apical spines on hind tibiae; hind tarsal plantulae; male subgenital plate; male suranal plate; female subgenital plate.

From our observations it is apparent that not all of these features are useful to characterise genera. The shape of male and female suranal and subgenital plates, for instance, used by both Ward (1997) and -RKQV &RRN LV VWURQJO\ VSHFLHV VSHFL¿F DQG ZKL WR GLIIHUHQWLDWH EHWZHHQ VSHFLHV LW LV WRR YDULDEOH generic level, with the possible exception of Pachyrhamma Brunner von Wattenwyl, 1888. The same can be said about body length. Even less useful are apical spines on the hind femora and the ventral OLQHDU VSLQHV RQ WKH KLQG IHPRUD WZR WUDLWV WKDW ZH Cook et al. 2010; Fitness et al. 2015).



Fig. 20. Tangle of Pleioplectron thomsoni (Chopard, 1923) comb. nov. on cave ceiling. Trotters Gorge, Otago.

While our observations mostly align with those of Ward (1997) and Johns & Cook (2013), there are nonetheless some differences. Ward (1997), for instance, described the female subgenital plate in Pleioplectron simplex Chopard, 1923 as "small, notched", with a matching illustration. In contrast, what we have observed (Fig. 10G) is in close agreement with the original description by Chopard (1923): "subgenital plate very small, almost hidden between the seventh sternite and the base of the ovipositor, rounded or little notched at the apex". Table 1 in Johns & Cook (2013) also seems to contain some inconsistencies, as Pleioplectron simplex and P. thomsoni comb. nov., for instance, are both described as having tarsal pads "bordered by spinules". We can clearly see these in the latter species, but not in the former. On the other hand, Johns & Cook (2013) reported the occasional presence of a retrolateral apical spine on the hind femur in P. thomsoni comb. nov., not in P. simplex or in Talitropsis sedilloti Bolivar, 1882. We have consistently found a retrolateral apical spine on the hind femur to be present in all three species. Johns & Cook's (2013) description of the dorsal spines on the hind tibiae of Pleioplectron simplex ("2 RU SDLUV LPPRYHDEOH VRFNHWHG VSLQHV ± SDLUV ¿[HG V count (20 to 33 prolateral, 23 to 35 retrolateral; see Table 1) or that of Richards (1959) (25–32 prolateral, 24–29 retrolateral). In the original description of Pleioplectron simplex, Hutton (1896) reported about 29 prolateral and 17 retrolateral dorsal spines on the hind tibiae. In agreement with Richards (1959), we assume the 17 to be a typographic error, 27 being the correct number, since the numbers of prolateral and UHWURODWHUDO GRUVDO VSLQHV DOZD\V FORVHO\ PDWFK HDF

While we note the inadequacy of the generic keys offered by Ward (1997) and by Johns & Cook (2013) in respect to Pleioplectron ZH FRQVLGHU LW SUHPDWXUH WR SUHVHQW DQ discussed in detail other genera of New Zealand Rhaphidophoridae.

A detail that has received little attention so far in the study of New Zealand Rhaphidophoridae is the structure of the antennae. The sensory hair tufts on the antennae of some (but not all) adult male Pleioplectron appear to be unique to this genus, and different from the antennal structures in Pachyrhamma Brunner von Wattenvyl, 1888 (Fea et al. 2019). While their precise function is currently unknown, typically, the sexually dimorphic structures of insect antennae have the purpose of increasing surface area for sensory perception in males (Fea et al. 2019).

Acknowledgements

Thanks to the New Zealand Department of Conservation for collecting permits and support of this biodiversity research. Rod Morris provided the information that enabled us to locate and collect specimens of Pleioplectron rodmorrisi sp. nov., and Warren Chinn provided additional information on the same species. Crystal Brindle offered hospitality and logistical support during many searches at Gouland Downs. Cor Vink and Jonathon Ridden of Canterbury Museum assisted us with access

Morgan-Richards M., Bulgarella M., Sivyer L., Dowle E.J., Hale M., McKean N.E. & Trewick S.A. 2017. Explaining large mitochondrial sequence differences within a population sample. Royal Society Open Science 4 (11): e170730. https://doi.org/10.1098/rsos.170730

Richards A.M. 1958. Revision of the Rhaphidophoridae (Orthoptera) of New Zealand Part I.—The Rhaphidophoridae of the Chatham Islands 1954 Expedition. Transactions of the Royal Society of New Zealand 85 (2): 263–274.

Richards A.M. 1959. Revision of the Rhaphidophoridae (Orthopetera) of New Zealand—Part V. The genus Pleioplectron Hutton, 1897. Transactions of the Royal Society of New Zealand 87 (3-4): 319–327.

Richards A.M. 1961a. Revision of the Rhaphidophoridae (Orthoptera) of New Zealand Part VIII. The species Turbottoplectron cavernae (Hutton) 1907. Transactions of the Royal Society of New Zealand, Zoology 1 (8): 113–119.

Richards A.M. 1961b. Revision of the Rhaphidophoridae (Orthoptera) of New Zealand Part IX. The genus Gymnoplectron Hutton, 1897. Transactions of the Royal Society of New Zealand, Zoology 1 (9): 323–326.

Simon C., Frati F., Beckenbach A., Crespi B.J., Liu H. & Flook P. 1994. Evolution, weighting, and phylogenetic utility of mitochondrial gene sequences and a compilation of conserved polymerase chain reaction primers. Annals of the Entomological Society of America 87: 651–701. https://doi.org/10.1093/aesa/87.6.651

Slatkin M. & Hudson R. 1991. Pairwise comparisons of mitochondrial DNA sequences in stable and exponentially growing populations. Genetics 129: 555–562.

Trewick S.A. & Morgan-Richards M. 2005. After the deluge: mitochondrial DNA indicates Miocene radiation and Pliocene adaptation of tree and giant weta (Orthoptera: Anostostomatidae). Journal of Biogeography 32: 295–309. https://doi.org/10.1111/j.1365-2699.2004.01179.x

Ward D.F. 1997. A new generic key to the New Zealand cave weta genera (Orthoptera: Rhaphidophoridae). New Zealand Natural Sciences 23: 13–17.

Manuscript received: 21 May 2019
Manuscript accepted: 21 August 2019
Published on: 12 November 2019
Topic editor: Gavin Broad
Desk editor: Danny Eibye-Jacobsen

Printed versions of all papers are also deposited in the libraries of the institutes that are members of the EJT consortium: Muséum national d'histoire naturelle, Paris, France; Meise Botanic Garden, Belgium; Royal Museum for Central Africa, Tervuren, Belgium; Royal Belgian Institute of Natural Sciences, Brussels, Belgium; Natural History Museum of Denmark, Copenhagen, Denmark; Naturalis Biodiversity Center, Leiden, the Netherlands; Museo Nacional de Ciencias Naturales-CSIC, Madrid, Spain; Real Jardín Botánico de Madrid CSIC, Spain; Zoological Research Museum Alexander Koenig, Bonn, Germany; National Museum, Prague, Czech Republic.

Errata corrigendum

In a previous paper published in the European Journal of Taxonomy, Fitness et al. 2018 (available from https://doi.org/10.5852/ejt.2018.468), we described and named a new species of Rhaphidophoridae, Miotopus richardsi. Since it is named after a female scientist, Aola Mary Richards, the correct declension is Miotopus richardsae, in compliance with article 31.1.2 of the ICZN Code.