

Fuchsia excorticata (Hemideina)

A A DA D. A. C, A A - A - C A D A D
B²

¹Ecology Group, Institute of Natural Resources, and ²Institute of Fundamental Sciences, Massey University, Private Bag 11-222, Palmerston North 4442, New Zealand

Abstract

Fuchsia excorticata (Hemideina) is a native New Zealand shrub. It is a highly invasive species, spreading rapidly across the country. The species is currently found in 78% of the country's area. *F. excorticata* is a highly invasive species, spreading rapidly across the country. The species is currently found in 78% of the country's area. *F. excorticata* is a highly invasive species, spreading rapidly across the country. The species is currently found in 78% of the country's area.

Key words: *Fuchsia excorticata*, Hemideina, New Zealand, invasive species.

INTRODUCTION

(Baker *et al.* 2006).

Handwritten musical notation on a staff, including notes, rests, and a treble clef.

Fuchsia excorticata (C. A. et al. 2008).
 (10) (B. 1995).
 500 600
 0.8 0.4 (& B. 1995). B.
F. excorticata (*H. novaeseelandiae*), (*Pr. novaeseelandiae*), (*A. melanura*), (*Z. lateralis*), (*Rhipidura fuliginosa*), (*Turdus merula*), (*Sturnus vulgaris*, C. et al. 1981; D. & D. 1994; B. 1995). et al. (2008).
F. excorticata
Fuchsia excorticata (et al. 2008).
 (*Hemideina*)
F. excorticata. (A. & 2005).
 (1992; 2006).
H. crassidens,
H. thoracica A. & 1995).
 (1978).

Husbandry of weta

Hemideina crassidens, *H. thoracica*
 15 × 15 × 9 10 × 10
 (1)

14 ± 1 C,
 (*Phormium tenax*),

***F. excorticata* seed survival**

F. excorticata
F. excorticata
F. excorticata
F. excorticata
 A.

F. excorticata
H. thoracica
H. crassidens

Table 1. *F. excorticata* seed survival in *Coriaria arborea* and *Eucalyptus* trees (n = 1000).

Host Tree	Survival (%)
<i>Coriaria arborea</i>	10%
<i>Eucalyptus</i>	85%

10% (0.10) × 0.22 = 0.022
 85% (0.85) × 0.22 = 0.187
 95% (0.95) × 0.22 = 0.209
 0.85 × 0.22 = 0.187
 (0.95 × 0.22) = 0.209
 B (1995): 54% *F. excorticata*
 et al. (2008)
 B (78%)
 et al. 2000).

(2007).

***F. excorticata* seed survival**

78% *F. excorticata*
 100%
 et al. 1998). A. (1995;
F. excorticata
 96 100%
 (B 1995), et al. (2008)
 (84%)
 D et al. (2006)

Movement of *F. excorticata* seeds

(C & C 1995). A.
 A.
 A.
 A.
 A.

F. excorticata

excorticata

REFERENCES

- B. (2005) A. *Tetranychus urticae*. *J. Appl. Entomol.* **129**, 173–5.
- B. (1991) *Keeping Wetas in Captivity: A Series of Nine Articles for Schools and Nature Lovers*. *NZ J. Ecol.* **29**, 430–6.
- B. (1994) *Q. Rev. Biol.* **69**, 31–51.
- B. (1998) *Biotropica* **30**, 150–61.
- B., A. & D. (2006) *New Phytol.* **172**, 412–28.
- B., C. (2006) *NZ J. Ecol.* **30**, 405–6.
- B., C. (1995) *Fuchsia excorticata*, *Griselinia littoralis*, *Macropiper excelsum*, *Melicytus ramiflorus*. *NZ J. Bot.* **33**, 131–40.
- C., C. A. & C. (1995) *Conserv. Biol.* **9**, 675–8.
- C., C. (2001) *Nature* **413**, 635–9.
- C., A. & D. (1981) *NZ J. Zool.* **8**, 87–91.
- C. & B. A. (1991) *Trends Ecol. Evol.* **6**, 193–5.
- D. C. (1869) *On the Origin of Species by Means of Natural Selection, Or the Preservation of Favoured Races in the Struggle for Life*. London: Chapman & Hall.
- D., C., & B. C. (2006) *Science* **311**, 1575.
- D., C. & B. C. (2001) *Hemideina femorata*, *Hemideina crassidens* (Blunell) (Insecta: Dermaptera). *The Biology of Wetas, King Crickets and Their Allies* (Ed. by D. C. & B. C.), pp. 243–57. CAB International, Wallingford.
- D., A. (1995) *Hypsipetes amaurotis*. *Res. Popul. Ecol.* **37**, 211–18.
- D., A., A. & D. (1998) *Ariolimax columbianus*. *Am. Midl. Nat.* **140**, 103–10.
- D., C. (2006) *Oikos* **112**, 481–92.
- D., C. & B. C. (1995) *Fuchsia*. *Ann. Mo. Bot. Gard.* **82**, 473–516.
- D., A. (2007) *Plants and Vegetation: Origins, Processes, Consequences* (Ed. by D. C.). pp. 365–79. Cambridge University Press, Cambridge.
- D., C. D. (2006) *NZ J. Ecol.* **30**, 273–8.
- D., A., A. & D. (2006) *Biological Invasions in New Zealand* (Ed. by D., A., A. & D.). pp. 227–45. Cambridge University Press, Cambridge.
- D., A., A. & D. (2009) *PLoS ONE* **4**, 5480.
- D., A., A. & D. (2004) *Austral Ecol.* **29**, 430–6.
- D., C., B. et al. (2010) *Ann. Bot.* **105**, 355–64.
- D., A. & D. (2008) *NZ J. Ecol.* **32**, 108–12.
- D., C. & D. (1994) *NZ J. Ecol.* **18**, 87–107.
- D., A. (1992) *Hemideina crassidens* (Blunell) (Insecta: Dermaptera). *NZ Entomol.* **15**, 1–8.
- D., C. & B. C. (1998) *Fuchsia excorticata*. *NZ J. Ecol.* **22**, 197–203.
- D., C. (1978) *NZ Entomol.* **6**, 400.